

HARMONIC DISCS OF SOLUTIONS TO THE COMPLEX HOMOGENEOUS MONGE-AMPÈRE EQUATION

JULIUS ROSS AND DAVID WITT NYSTRÖM

ABSTRACT. We study regularity properties of solutions to the Dirichlet problem for the complex Homogeneous Monge-Ampère equation. We show that for certain boundary data on \mathbb{P}^1 the solution Φ to this Dirichlet problem is connected via a Legendre transform to an associated flow in the complex plane called the Hele-Shaw flow. Using this we determine precisely the harmonic discs associated to Φ . We then give examples for which these discs are not dense in the product, and also prove that this situation persists after small perturbations of the boundary data.

Let (X, ω) be a compact Kähler manifold of dimension n and $\mathbb{D} \subset \mathbb{C}$ be the open unit disc. Consider boundary data consisting of a family $\omega + dd^c \phi(\cdot, \tau)$ of Kähler forms where $\phi(\cdot, \tau)$ is a smooth function on X for $\tau \in \partial\mathbb{D}$. The Dirichlet problem for the complex Homogeneous Monge Ampère equation (HMAE) asks for a function Φ on $X \times \overline{\mathbb{D}}$ such that

$$\begin{aligned}\Phi(\cdot, \tau) &= \phi(\cdot, \tau) \text{ for } \tau \in \partial\mathbb{D}, \\ \pi^* \omega + dd^c \Phi &\geq 0, \\ (\pi^* \omega + dd^c \Phi)^{n+1} &= 0.\end{aligned}$$

We say Φ is a *regular solution* if it is smooth and $\omega + dd^c \Phi(\cdot, \tau)$ is a Kähler form for all $\tau \in \overline{\mathbb{D}}$. By an example of Donaldson [12] we know there exist smooth boundary data for which there does not exist a regular solution. Nevertheless, the equation always has a unique weak solution, which by the work of Chen [6] with complements by Błocki [5] we know is at least “almost” $C^{1,1}$ (so in particular $C^{1,\alpha}$ for any $\alpha < 1$). See [14] for a recent survey.

A more subtle aspect of the regularity of solutions to the HMAE is the question of existence and distribution of harmonic discs.

Definition. Let $g: \mathbb{D} \rightarrow X$ be holomorphic. We say that the graph of g is a *harmonic disc* (with respect to Φ) if Φ is $\pi^* \omega$ -harmonic (i.e. $\pi^* \omega + dd^c \Phi$ vanishes) along this graph.

As is well known, a regular solution to the HMAE yields a complex foliation of $X \times \overline{\mathbb{D}}$ whose leaves restrict to harmonic discs in $X \times \mathbb{D}$. Even when the solution is not regular, the existence of such harmonic discs is important; for instance along such a harmonic disc the density of the varying measure $\omega_{\phi(\cdot, \tau)}^n$ is essentially log-subharmonic (see [1] [7] [4, Sec 3.2]).

It was hoped that any weak solution would enjoy a weaker form of regularity, so that a dense open subset of $X \times \mathbb{D}$ would be foliated by harmonic disc, but as we will see this is not always the case.

During this work JR was supported by an EPSRC Career Acceleration Fellowship (EP/J002062/1). DWN has received funding from the People Programme (Marie Curie Actions) of the European Union’s Seventh Framework Programme (FP7/2007-2013) under REA grant agreement no 329070.

This paper describes a correspondence between on the one hand the HMAE when $X = \mathbb{P}^1$ and the boundary data has a certain kind of symmetry and on the other hand the so-called “Hele-Shaw” flow in the plane. As a result we see that the set of harmonic discs is determined by the topology of the flow.

To state precise results, let ω_{FS} denote the Fubini-Study form on \mathbb{P}^1 and ϕ be a smooth Kähler potential, i.e. a smooth function on \mathbb{P}^1 such that $\omega_{FS} + dd^c\phi$ is Kähler. Let ρ denote the usual \mathbb{C}^\times -action on \mathbb{P}^1 which acts by multiplication on $\mathbb{C} \subset \mathbb{P}^1$. We consider the function $\phi(z, \tau) := \phi(\rho(\tau)z)$ as boundary data over $\mathbb{P}^1 \times \partial\mathbb{D}$, so for each $\tau \in \partial\mathbb{D}$ we have a Kähler form $\omega_{FS} + dd^c\phi(\cdot, \tau)$. We show that the solution Φ to the Homogeneous Monge-Ampère equation (see 2.1 for the definition) with this boundary data is intimately connected to the Hele-Shaw flow

$$\Omega_t := \{z : \psi_t(z) < \phi(z)\}$$

where

$$\psi_t := \sup\{\psi : \psi \text{ is usc and } \psi \leq \phi \text{ and } \omega_{FS} + dd^c\psi \geq 0 \text{ and } \nu_0(\psi) \geq t\}.$$

By this we mean the supremum is over all upper semicontinuous (usc) functions from \mathbb{P}^1 to $\mathbb{R} \cup \{-\infty\}$ with these properties, and $\nu_0(\psi)$ denotes the order of the logarithmic singularity (Lelong number) of ψ at $0 \in \mathbb{C} \subset \mathbb{P}^1$. In fact we show that the solution Φ and the family ψ_t are related via a Legendre transform.

Using this we prove the following:

Theorem 1. *Let Φ be the solution to the HMAE with boundary data ϕ and $g : \mathbb{D} \rightarrow \mathbb{P}^1$ be holomorphic. Then the graph of g is a harmonic disc of Φ if and only if either*

- (1) $g \equiv 0$, or
- (2) $g(\tau) = \tau^{-1}z$ for some fixed $z \in \mathbb{P}^1 \setminus \Omega_1$, or
- (3) $\tau \mapsto \tau g(\tau)$ is a Riemann mapping for a simply connected Hele-Shaw domain Ω_t that maps $0 \in \mathbb{D}$ to $0 \in \Omega_t$.

Remark. In [20] the authors prove that Ω_t is simply connected for $0 < t \ll 1$, so there is always an infinite number of harmonic discs of the form (3).

The Hele-Shaw flow Ω_t has a physical interpretation as describing the expansion of a liquid in a medium with permeability inversely proportional to $\Delta(\phi + \ln(1 + |z|^2))$. Guided by this one can rather easily find potentials ϕ for which at some time t the flow domain Ω_t becomes multiply connected as in Figure 1.




FIGURE 1. The Hele-Shaw flow developing a multiply connected domain

This then translates into an obstruction to the presence of harmonic discs of the associated solution to the HMAE:

Theorem 2. *There exist smooth boundary data $\phi(\cdot, \tau)$ for which the solution to the Dirichlet problem for the HMAE has the following property: there exists an open set U in $\mathbb{P}^1 \times \mathbb{D}$ meeting $\mathbb{P}^1 \times \partial\mathbb{D}$, such that no harmonic disc intersects U .*

Next we address the question whether generic boundary data give rise to solutions with a weak form of regularity. The following theorem answers that question negatively.

Theorem 3. *There exist smooth boundary data $\phi(\cdot, \tau)$ for which the following is true: there exist a nonempty open set U' in $\mathbb{P}^1 \times \mathbb{D}$ and an $\epsilon > 0$ such that if $\phi'(z, \tau)$ is any smooth boundary data with*

$$\|\phi' - \phi\|_{C^2(\mathbb{P}^1 \times \partial\mathbb{D})} < \epsilon$$

and Φ' is the associated solution to the HMAE then no harmonic disc (associated to Φ') passes through U' .

The first theorem gives solutions that are not “partially smooth” and the second examples whose perturbed solutions are not “almost smooth”, in apparent contradiction with [7, Thm. 1.3.2] and [7, Thm. 1.3.4] respectively (see Section 6).

Comparison with other works. In this paper we consider the Hele-Shaw flow with varying permeability starting from the origin which has been considered before by Hedenmalm-Shimorin; in fact the basic properties of the flow that we use are a small variant of those in [16]. There is a much larger literature on the Hele-Shaw flow (which usually considers the case of constant permeability but with non-trivial initial condition Ω_0) for which we refer the reader to the book [15] of Gustafsson-Vasil’ev and the references therein.

In [12] Donaldson gave the first example of boundary data for which the Dirichlet problem for the HMAE on some $X \times \overline{\mathbb{D}}$ has no regular solution.

One motivation for studying the regularity of solutions to the HMAE is the work of Semmes [21] and Donaldson [12], which shows that the geodesic equation in the space of Kähler metrics on a compact Kähler manifold (X, ω) cohomologous to ω can be cast as a Dirichlet problem for the HMAE on $X \times A$ where A is an annulus. In this way the question of geodesic connectivity is translated into a question of regularity for solutions to the HMAE. In [18] Lempert-Vivas found such boundary data for which the solution failed to be regular, and thus showed that not all pairs of cohomologous Kähler metrics can be connected by a geodesic (see also later work by Darvas-Lempert [10] and Darvas [11]).

In [10] examples are given for which the solution (again with the base being an annulus rather than a disc) does not have continuous second derivative. We expect that the examples considered here in fact fail to be twice differentiable at any point $(z, 1)$ where $z \in \mathbb{P}^1$ is a self-intersection point of the boundary of a simply connected Hele-Shaw domain Ω_t . Finally, we refer the reader to [14, Chapter 2] for a discussion of the analogous problem of the complex HMAE for domains in \mathbb{C}^n .

The connection between the HMAE and the Hele-Shaw flow has been studied previously by the authors in [20] (see also [13] for another connection between the HMAE and free boundary problems). The results there are in the opposite direction to those here, in that we use known regularity results of the HMAE (and thus the existence of the associated foliation by holomorphic discs) to prove short time regularity of the Hele-Shaw flow.

Acknowledgements. We would particularly like to thank Bo Berndtsson for his close reading of the first version of this paper which has led to many improvements. We also thank Robert Berman, Zbigniew Błocki, Laszlo Lempert and Yanir Rubinstein for discussions concerning this work.

1. THE HELE-SHAW FLOW

1.1. Definition and Basic Properties. Assume ϕ is a smooth Kähler potential on \mathbb{P}^1 , so $\omega_\phi := \omega_{FS} + dd^c\phi$ is Kähler. Given $t \in [0, 1]$ we define

$$\psi_t = \sup\{\psi : \psi \text{ is usc and } \omega_\psi \geq 0 \text{ and } \psi \leq \phi \text{ and } \nu_0(\psi) \geq t\}. \quad (1)$$

Here $\omega_\psi := \omega_{FS} + dd^c\psi$ and ν_0 denotes the Lelong number, so $\nu_0(\psi) \geq t$ means that there is a constant C such that $\psi(z) \leq t \ln|z|^2 + C$ for all z near $0 \in \mathbb{C} \subset \mathbb{P}^1$. As the usc regularization of ψ_t is itself a candidate for the envelope we see that ψ_t is usc.

We now define the Hele-Shaw flow by

$$\Omega_t := \Omega_t^\phi := \{z \in \mathbb{P}^1 : \psi_t(z) < \phi(z)\}. \quad (2)$$

Proposition 1.1. (*Basic Properties of Hele-Shaw flow*)

- (1) $\Omega_0 = \emptyset$ and $0 \in \Omega_t$ for $t > 0$.
- (2) Ω_t is open, connected and $\partial\Omega_t$ has measure zero.
- (3) ψ_t is $C^{1,1}$ on $\mathbb{P}^1 \setminus \{0\}$.
- (4)

$$\omega_{\psi_t} = (1 - \chi_{\Omega_t})\omega_\phi + t\delta_0$$

in the sense of currents. Here χ_A denotes the characteristic function of a set A , and δ_0 the Dirac delta.

(5)

$$\text{Area}(\Omega_t) := \int_{\Omega_t} \omega_\phi = t.$$

Proof. This techniques used for this are standard (see e.g. [16]) but for the convenience of the reader we sketch some details.

Clearly $\psi_0 \equiv \phi$ and thus $\Omega_0 = \emptyset$ so the Lemma is obviously true for $t = 0$. Thus let $t > 0$. That $0 \in \Omega_t$ is obvious. Since ψ_t is upper semicontinuous $\psi_t < \phi$ in a neighbourhood of the origin. It follows from standard potential theory that $\psi_t + \ln(1 + |z|^2)$ is harmonic in any open set contained in $\Omega_t \setminus \{0\}$, so in particular ψ_t is smooth on any punctured disc D centered at the origin contained in Ω_t . Define

$$\psi' = \sup \left\{ \begin{array}{l} \psi \text{ is usc on } \mathbb{P}^1 \setminus D \text{ with } \omega_{FS|_{\mathbb{P}^1 \setminus D}} + dd^c\psi \geq 0 \text{ and } \psi \leq \phi|_{\mathbb{P}^1 \setminus D} \\ \text{and } \psi \leq \psi_t \text{ on } \partial D \end{array} \right\}.$$

Since $\psi_t|_{\mathbb{P}^1 \setminus D}$ is a candidate for the envelope ψ' we get that $\psi' \geq \psi_t|_{\mathbb{P}^1 \setminus D}$. On the other hand, if $\epsilon > 0$ then extending $\max(\psi_t, \psi' - \epsilon)$ by ψ_t on D gives an ω_{FS} -subharmonic function on \mathbb{P}^1 which is thus a candidate for the envelope defining ψ_t . Hence $\psi' \leq \psi_t|_{\mathbb{P}^1 \setminus D} + \epsilon$ and so $\psi' = \psi_t|_{\mathbb{P}^1 \setminus D}$.

If we let $w = 1/z$ then $\psi'(w) + \ln(1 + |w|^2)$ now solves a standard free boundary problem on the disc $\mathbb{P}^1 \setminus D$ with obstacle given by $\phi(w) + \ln(1 + |w|^2)$ and boundary condition given by $\psi_t(w) + \ln(1 + |w|^2)$ restricted to ∂D . So (3) follows from standard theory of free boundary problems (e.g. [9, Thm 2.3]) and (4) follows from (3) on $\mathbb{P}^1 \setminus D$. On the other hand the function $t \ln|z|^2 + C$ for some constant C is a candidate for the envelope defining ψ_t , so $\nu_0(\psi_t) = t$, giving (4) on all of \mathbb{P}^1 . Moreover (5) in turn follows from (4) by Stokes theorem.

That Ω_t is open of course also follows from ψ_t being continuous. The proof that Ω_t is connected is as in [16, Prop 2.6]. Finally the fact that $\partial\Omega_t$ has zero measure again follows from standard theory of free boundary problems, see e.g [8, p296]. In fact even more is proved in [8], namely that each component of the boundary consists of a finite number of rectifiable Jordan curves. \square

Remark 1.2. Item (4) in particular implies that the sets Ω_t together with ω_ϕ contain the same information as the functions ψ_t , and so we also think of this latter collection as the Hele-Shaw flow. Items (3), (4) and (5) of the above Proposition also follow from more general work of Berman [2, Sec 4].

Clearly by definition if $t \leq t'$ then $\Omega_t \subset \Omega_{t'}$ so this is an increasing flow of subsets of \mathbb{P}^1 . Observe also that if ϕ is rotation invariant then so is the flow Ω_t .

An important fact is that the family of functions ψ_t is concave in t .

Proposition 1.3. *For any given z we have that $\psi_t(z)$ is concave, decreasing and continuous in t for $t \in [0, 1]$.*

Proof. Let us define $\psi_t := \phi$ for $t < 0$. It is then clear that ψ_t is concave in t since if $t = at_1 + (1-a)t_2$ where $a \in [0, 1]$ and $t_1, t_2 \in (-\infty, 1]$ then

$$a\psi_{t_1} + (1-a)\psi_{t_2} \leq \psi_t$$

simply because the LHS has at least Lelong number t at the origin while being bounded from above by ϕ . That it is decreasing is obvious. That ψ_t decreases with t then implies that $\lim_{t \rightarrow s-} \psi_t$ is ω_{FS} -subharmonic and thus one sees that

$$\lim_{t \rightarrow s-} \psi_t = \psi_s,$$

i.e. ψ_t is left-continuous in t . This combined with concavity implies continuity. \square

1.2. Multiply Connected Hele-Shaw Domains. As already mentioned, the sets Ω_t have a physical interpretation. They describe the flow obtained by injecting a fluid at a point between two parallel plates between which there is a medium with permeability inversely proportional to $\Delta(\phi + \ln(1 + |z|^2))$. As such it is intuitively clear that there will be ϕ for which this flow becomes multiply connected. In fact suppose we arrange so $\Delta(\phi + \ln(1 + |z|^2))$ is very small on some Jordan curve going through the origin, while $\Delta(\phi + \ln(1 + |z|^2))$ being relatively large in two regions separated by the curve. The the flow will then cover the curve before having the chance to engulf either of the regions, thereby giving rise to multiply connected Hele-Shaw domains. We now prove that this does indeed happen for suitable choices of ϕ .

Proposition 1.4. *There exists a smooth Kähler potential on \mathbb{P}^1 whose associated Hele-Shaw flow has the following property: there exist two times $0 < t_1 < t_2 < 1$ such that for all $t \in (t_1, t_2)$ the Hele-Shaw domain Ω_t is not simply connected.*

Proof. We will construct a ϕ with the following property: there exist two times $0 < t_1 < t_2 < 1$, a Jordan curve γ which passes through the origin and two points p, q on opposite sides of γ such that Ω_{t_1} contains γ while neither p or q lie in Ω_{t_2} . By monotonicity of the flow, it follows immediately that for any $t \in (t_1, t_2)$ the Hele-Shaw domain Ω_t is multiply connected.

So pick a Jordan curve γ in \mathbb{P}^1 which passes through the origin, and let U_1 and U_2 denote the two connected components of the complement of γ . Let f be a smooth nonnegative function on \mathbb{P}^1 which is zero in a neighbourhood U_γ of γ and such that

$$\int_{U_1} f \omega_{FS} = \int_{U_2} f \omega_{FS} = 1/2.$$

Let ϕ_f be a smooth function such that

$$\omega_{\phi_f} = f \omega_{FS}.$$

Fix

$$0 < t_0 < 1/4,$$

and set

$$\psi_f := \sup\{\psi : \omega_\psi \geq 0, \psi \leq \phi_f, \nu_0(\gamma) \geq t_0\}.$$

So $\omega_{\psi_f} = \chi_{\{\psi_f = \phi_f\}} \omega_{\phi_f} + t_0 \delta_0$ and we conclude that $\psi_f - \phi_f$ is harmonic in $U_\gamma \setminus \{0\}$ while having a singularity at the origin. Clearly $\psi_f - \phi_f \leq 0$ on U_γ . Thus we conclude from the maximum principle that in fact $\psi_f - \phi_f < 0$ in U_γ .

Now let

$$\psi_\epsilon := \sup\{\psi : \omega_\psi \geq -\epsilon \omega_{FS}, \psi \leq \phi_f, \nu_0(\psi) \geq t_0\}.$$

Then $\psi_\epsilon - \phi_f$ decreases to $\psi_f - \phi_f$ as ϵ tends to zero (as for any decreasing sequence of subharmonic functions the limit is also subharmonic, and thus a candidate for $\psi_f - \phi_f$). In particular if z is a fixed point on γ then for ϵ sufficiently small $\psi_\epsilon - \phi_f$ is strictly negative at z , and hence by upper semicontinuity of $\psi_\epsilon - \phi_f$ this is true in a neighbourhood of z . Thus for $\epsilon \ll 1$ sufficiently small, $\psi_\epsilon - \phi_f$ is negative on the compact set γ .

Pick such a small ϵ and let

$$\phi := \frac{\phi_f}{1 + \epsilon}.$$

Then $\omega_\phi = (1 + \epsilon)^{-1}(\epsilon \omega_{FS} + \omega_{\phi_f})$ so ϕ is a Kähler potential. By construction it now follows that if we let $t_1 := (1 + \epsilon)^{-1}t_0$ then $\gamma \subset \Omega_{t_1}^\phi$. Observe that $0 < t_1 < 1/4$.

We have that

$$\text{Area}_\phi(U_1) := \int_{U_1} \omega_\phi = \int_{U_1} \omega_{FS} + \frac{dd^c \phi_f}{1 + \epsilon} \geq \frac{1}{1 + \epsilon} \int_{U_1} \omega_{\phi_f} > 1/4,$$

and similarly $\text{Area}_\phi(U_2) > 1/4$. Now if we pick some t_2 such that $0 < t_1 < t_2 < 1/4$, then

$$\text{Area}_\phi(\Omega_{t_2}^\phi) = t_2 < 1/4 < \text{Area}_\phi(U_1)$$

and similarly

$$\text{Area}_\phi(\Omega_{t_2}^\phi) < \text{Area}_\phi(U_2).$$

In particular the sets $U_1 \setminus \Omega_{t_2}^\phi$ and $U_2 \setminus \Omega_{t_2}^\phi$ must both be nonempty, which allows us to pick points p and q in the complement of $\Omega_{t_2}^\phi$ on either side of γ . This then concludes the proof. \square

2. THE LEGENDRE TRANSFORM BETWEEN THE HMAE AND THE HELE-SHAW FLOW

We shall focus on a simple case of the complex Homogeneous Monge Ampère equation. Suppose that $\phi(\cdot, \tau)$ is a smooth family of Kähler potentials parameterized by $\tau \in \partial\mathbb{D}$. We denote by π the projection $\mathbb{P}^1 \times \overline{\mathbb{D}} \rightarrow \mathbb{P}^1$.

Definition 2.1. The *solution* Φ to the Homogeneous Monge-Ampère equation with boundary data $\phi(\cdot, \tau)$ is the function on $\mathbb{P}^1 \times \overline{\mathbb{D}}$ given by

$$\Phi = \sup\{\psi : \psi \text{ is usc and } \pi^* \omega_{FS} + dd^c \psi \geq 0 \text{ and } \psi(\cdot, \tau) \leq \phi(\cdot, \tau) \text{ for } \tau \in \partial\mathbb{D}\}.$$

From general theory $\pi^* \omega_{FS} + dd^c \Phi \geq 0$ as a current, has the boundary value $\Phi(\cdot, \tau) = \phi(\cdot, \tau)$ for $\tau \in \partial\mathbb{D}$ and solves the equation

$$(\pi^* \omega_{FS} + dd^c \Phi)^2 = 0$$

in the sense of Bedford-Taylor. As is well known, due to Chen [6] with complements by Błocki [5], the solution is almost $C^{1,1}$, and in fact since in our case $(\mathbb{P}^1, \omega_{FS})$ has nonnegative sectional curvature the solution is truly $C^{1,1}$ by the result of Błocki [5, Thm.

1.4] (see also the recent work of Berman [3] for a proof the weak solution is $C_{loc}^{1,1}$ along the original lines of Bedford-Taylor).

Recall ρ denotes the usual \mathbb{C}^\times -action on \mathbb{P}^1 which acts by multiplication on $\mathbb{C} \subset \mathbb{P}^1$. Letting ϕ be a Kähler potential as before, we wish to consider the function

$$\phi(z, \tau) := \phi(\rho(\tau)z)$$

as boundary data to the HMAE on $\mathbb{P}^1 \times \overline{\mathbb{D}}$. Let Φ denote the weak solution to the corresponding HMAE as in (2.1). The goal in this section is to show that Φ is connected via a Legendre transform to the Hele-Shaw flow on \mathbb{P}^1 taken with respect to ϕ .

To do this, consider the envelope on $\mathbb{P}^1 \times \overline{\mathbb{D}}$ given by

$$\tilde{\Phi} = \sup\{\psi : \psi \text{ is usc, } \pi^*\omega_{FS} + dd^c\psi \geq 0, \psi(z, \tau) \leq \phi(z) \text{ for } \tau \in \partial\mathbb{D} \text{ and } \nu_{(0,0)}(\psi) \geq 1\}$$

(so the boundary data is independent of τ , and the ψ have Lelong number at least one at the point $(0, 0)$). Then by standard arguments, $\tilde{\Phi}$ is usc, $\pi^*\omega_{FS} + dd^c\tilde{\Phi} \geq 0$ and $(\pi^*\omega_{FS} + dd^c\tilde{\Phi})^2 = 0$ away from $(0, 0)$.

Remark 2.2. The function $\tilde{\Phi}$ solves the HMAE over the punctured disc $\overline{\mathbb{D}}^\times$ with boundary data independent of τ , and thus is a weak geodesic ray emanating from ϕ .

Proposition 2.3. *We have that*

$$\Phi(z, \tau) + \ln|\tau|^2 + \ln(1 + |z|^2) = \tilde{\Phi}(\tau z, \tau) + \ln(1 + |\tau z|^2) \text{ for } (z, \tau) \in \mathbb{P}^1 \times \overline{\mathbb{D}}^\times.$$

Proof. Consider the space \mathcal{X} we get by blowing up $\mathbb{P}^1 \times \overline{\mathbb{D}}$ at the point $(\infty, 0)$. Let μ_1 denote the modification map, and let E_1 denote the exceptional divisor. The central fibre thus consists of two copies of \mathbb{P}^1 . We call the other one E_2 , and if we blow down this one we get again $\mathbb{P}^1 \times \overline{\mathbb{D}}$. Call this modification map μ_2 , so E_2 is now the exceptional divisor of μ_2 . One can now check that

$$\mu_2 \circ \mu_1^{-1}(z, \tau) = (\tau z, \tau),$$

except for being undefined at the point $(\infty, 0)$.

We have two ways to pull back the Fubini-Study form to \mathcal{X} , via μ_1 or μ_2 , and we want to describe how they are related. For this let f denote the function on \mathcal{X} defined as

$$f := (\ln(|\tau|^2 + |z|^2) - \ln(1 + |z|^2)) \circ \mu_2.$$

Then a simple calculation shows that

$$dd^c f = \mu_1^*(\pi^*\omega_{FS}) - (\mu_2^*(\pi^*\omega_{FS}) - [E_2]),$$

where $[E_2]$ denotes the current of integration along E_2 . This then shows that

$$\Phi \circ \mu_1 \circ \mu_2^{-1} + \ln(|\tau|^2 + |z|^2) - \ln(1 + |z|^2)$$

is a candidate for the envelope defining $\tilde{\Phi}$, while reversely

$$\tilde{\Phi} \circ \mu_2 \circ \mu_1^{-1} + f \circ \mu_1^{-1}$$

is a candidate for the envelope defining Φ . It follows that

$$\Phi \circ \mu_1 + f = \tilde{\Phi} \circ \mu_2$$

and we simply get the formula of the Proposition by first taking μ_1^{-1} . \square

Remark 2.4. From the above we see that

$$\Phi(z, \tau) - \tilde{\Phi}(\tau z, \tau) = \ln \left(\frac{1 + |\tau z|^2}{|\tau|^2(1 + |z|^2)} \right)$$

which is smooth on $\mathbb{P}^1 \times \overline{\mathbb{D}}^\times$. Hence as Φ is $C^{1,1}$ on $\mathbb{P}^1 \times \overline{\mathbb{D}}^\times$, the same is true of $\tilde{\Phi}$.

Lemma 2.5.

$$\tilde{\Phi}(z, \tau) \leq \ln(|\tau|^2 + |z|^2) - \ln(1 + |z|^2) + \max(\phi). \quad (3)$$

Proof. First assume that ϕ is identically zero. Then clearly Φ is also identically zero so from Proposition 2.3 we see that

$$\tilde{\Phi}(z, \tau) = \ln(|\tau|^2 + |z|^2) - \ln(1 + |z|^2).$$

The lemma then follows from the obvious monotonicity property of $\tilde{\Phi}$. \square

Now we wish to connect $\tilde{\Phi}$ (and hence Φ) with the Hele-Shaw flow of ϕ . For any $t \in \mathbb{R}$ we let

$$\psi_t = \sup\{\psi : \psi \text{ is usc and } \omega_\psi \geq 0 \text{ and } \psi \leq \phi \text{ and } \nu_0(\psi) \geq t\}. \quad (4)$$

Note that for $t < 0$, $\psi_t = \phi$ while if $t > 1$ we get that $\psi_t \equiv -\infty$. For $t \in [0, 1]$ we recognize ψ_t as the envelopes that contain the same data as the Hele-Shaw flow $\Omega_t = \{\psi_t < \phi\}$.

We saw in Section 1 that for a fixed z the function $\psi_t(z)$ is concave in t . On the other hand, the function $\tilde{\Phi}(z, e^{-s/2})$ is subharmonic and independent of the imaginary part of s and thus $\tilde{\Phi}(z, e^{-s/2})$ is convex in s (where we now think of s as a real variable taking values in $[0, \infty)$). We can then define $\tilde{\Phi}(z, e^{-s/2})$ to be $+\infty$ for $s < 0$ to get a convex function defined on the whole of \mathbb{R} .

Now we recall the definition of the (one-variable) Legendre transform (or convex conjugate).

Definition 2.6. Given a function $u : \mathbb{R} \rightarrow \mathbb{R} \cup \{+\infty\}$ the Legendre transform $\hat{u} : \mathbb{R} \rightarrow \mathbb{R} \cup \{+\infty\}$ is the convex function defined as

$$\hat{u}(y) := \sup_{x \in \mathbb{R}^n} \{xy - u(x)\}.$$

The Fenchel-Moreau Theorem (see e.g. [19]) now asserts that the Legendre transform is an involution (i.e. $\hat{\hat{u}} = u$) precisely on the set of convex lower semicontinuous functions.

The next Theorem says that $u(s) := \tilde{\Phi}(z, e^{-s/2}) + s$ is the Legendre transform of $-\psi_t(z)$ and vice versa.

Theorem 2.7.

$$\psi_t(z) = \inf_{|\tau| > 0} \{\tilde{\Phi}(z, \tau) - (1 - t) \ln |\tau|^2\} \quad (5)$$

and

$$\tilde{\Phi}(z, \tau) = \sup_t \{\psi_t(z) + (1 - t) \ln |\tau|^2\}. \quad (6)$$

Proof. First we note that

$$\psi_t(z) + (1 - t) \ln |\tau|^2$$

is a candidate for the envelope defining $\tilde{\Phi}$ so by definition we get that the LHS is less than or equal to the RHS in (5). Since $\tilde{\Phi}(z, \tau)$ is independent of the argument of τ it follows from Kiselman's minimum principle (see [17]) that the RHS defines an ω_{FS} -sh function on \mathbb{P}^1 , which we will denote by $\tilde{\psi}_t$. Clearly $\tilde{\psi}_t \leq \phi$, so if we can show that it has logarithmic singularity of order at least t at the origin, then it would follow from the

definition of ψ_t as the supremum of all such functions that $\tilde{\psi}_t \leq \psi_t$. Since for a fixed z the function $\tilde{\Phi}(z, e^{-s/2})$ is subharmonic and independent of the imaginary part of s we get that $\tilde{\Phi}(z, e^{-s/2})$ is convex in s (where we now think of s as a real variable taking values in $[0, \infty)$). Thus for a fixed z

$$-\tilde{\psi}_t(z) = \inf_{s \geq 0} \{ \tilde{\Phi}(z, e^{-s/2}) + (1-t)s \} = \sup_{s \in \mathbb{R}} \{ ts - (\tilde{\Phi}(z, e^{-s/2}) + s) \}$$

is the Legendre transform of a convex function. Using the inequality (3) we get that

$$\tilde{\psi}_t(z) \leq \inf_{s \geq 0} \{ \ln(e^{-s} + |z|^2) + (1-t)s \} - \ln(1 + |z|^2) + \max(\phi).$$

By elementary means one easily checks that if $t \in (0, 1)$ then

$$\inf_{s \geq 0} \{ \ln(e^{-s} + |z|^2) + (1-t)s \} = t(\ln |z|^2 - \ln t) - (1-t) \ln(1-t)$$

which shows that indeed $\tilde{\psi}_t$ has a logarithmic singularity of order t , at least when $t \in (0, 1)$. For $t = 0$ one notes that $\phi(z) + \ln |\tau|^2$ is a candidate for the envelope defining $\tilde{\Phi}$, so $\phi(z) + \ln |\tau|^2 \leq \tilde{\Phi}$, which implies that $\tilde{\psi}_0 = \phi$. For $t = 1$, the fact that

$$\tilde{\psi}_1(z) \leq \tilde{\Phi}(z, 0)$$

implies that it has the right singularity as well. The case that $t < 0$ and $t > 1$ are immediate, so this proves (5).

We thus see that

$$-\psi_t(z) = \sup_{s \in \mathbb{R}} \{ ts - (\tilde{\Phi}(z, e^{-s/2}) + s) \},$$

i.e. that $-\psi_t(z)$ is the Legendre transform of $u(s) := \tilde{\Phi}(z, e^{-s/2}) + s$. We also know that $u(s)$ is convex and lower semicontinuous (since it is continuous on $[0, \infty)$ and constantly $-\infty$ on $(-\infty, 0)$) so by the Fenchel-Moreau Theorem we get that $u(s)$ is the Legendre transform of $-\psi_t(z)$. This is exactly what is asserted in (6). \square

2.1. The Hamiltonian. We will have use for the function H on $\mathbb{P}^1 \times \overline{\mathbb{D}}^\times$ defined as

$$H(z, \tau) := \frac{\partial}{\partial s} \tilde{\Phi}(z, e^{-s/2}),$$

where $s := -\ln |\tau|^2$ (when $|\tau| = 1$ and thus $s = 0$ we take the right derivative). As $\tilde{\Phi}$ is $C^{1,1}$ on $\mathbb{P}^1 \times \overline{\mathbb{D}}^\times$ the function H is well-defined and continuous (even Lipschitz but we will not need this).

Proposition 2.8.

$$H(z, 1) + 1 = \sup \{ t : \psi_t(z) = \phi(z) \} = \sup \{ t : z \notin \Omega_t \}.$$

Proof. From (6) we see that if $\psi_t(z) = \phi(z)$ then

$$\tilde{\Phi}(z, e^{-s/2}) \geq (t-1)s + \phi(z)$$

and thus

$$H(z, 1) \geq \sup \{ t : \psi_t(z) = \phi(z) \} - 1.$$

Recall that for a fixed z the function $\psi_t(z)$ is concave and decreasing in t . Thus if $\psi_t(z) \leq a < 0$ then it follows from (6) that

$$\tilde{\Phi}(z, e^{-s/2}) \leq \max((t-1)s, -a) + \phi(z)$$

and so $H(z, 1) \leq t - 1$, which proves the proposition. \square

Proposition 2.9. *For $0 < |\tau| < 1$ we have that*

$$H(z, \tau) = t - 1 \iff \tilde{\Phi}(z, \tau) = \psi_t(z) - (1 - t) \ln |\tau|^2.$$

Proof. Fix a point (z_0, τ_0) , $0 < |\tau_0| < 1$. From (6) we see that

$$\tilde{\Phi}(z_0, \tau_0) = \sup_{t \in [0, 1]} \{\psi_t(z_0) + (1 - t) \ln |\tau_0|^2\}.$$

Since $\psi_t(z_0)$ is continuous in t we must have that $\tilde{\Phi}(z_0, \tau_0) = \psi_{t_0}(z_0) + (1 - t_0) \ln |\tau_0|^2$ for some $t_0 \in [0, 1]$. Since we always have that

$$\tilde{\Phi}(z, e^{-s/2}) \geq \psi_{t_0}(z) - (1 - t_0)s$$

it follows that

$$H(z_0, \tau_0) = \frac{\partial}{\partial s} \Big|_{s = -\ln |\tau_0|^2} (\psi_{t_0}(z) - (1 - t_0)s) = t_0 - 1.$$

□

3. HARMONIC DISCS

As above, let Φ be the weak solution to the HMAE with boundary data $\phi(z, \tau) = \phi(\rho(\tau)z)$. Recall that if $g: \mathbb{D} \rightarrow \mathbb{P}^1$ is holomorphic then we say the graph of g is a harmonic disc if Φ is $\pi^* \omega_{FS}$ -harmonic along this graph.

Theorem 3.1. *Let $g: \mathbb{D} \rightarrow \mathbb{P}^1$ be holomorphic. Then the graph of g is a harmonic disc of Φ if and only if either (1) $g \equiv 0$ or (2) $g(\tau) = \tau^{-1}z$ where $z \in \Omega_1^c$ or (3) $\tau \mapsto \tau g(\tau)$ is a Riemann mapping for a simply connected Hele-Shaw domain Ω_t that maps $0 \in \mathbb{D}$ to $0 \in \Omega_t$. The function H is constant along the associated discs $\{(\tau g(\tau), \tau)\}$, in the first case $H = -1$, in the second case $H = 0$ while in the third case $H = t - 1$.*

Lemma 3.2. *If $f: D_1 \rightarrow D_2$ is a proper holomorphic map between two open domains in \mathbb{P}^1 then the number of preimages $N_p := \#\{f^{-1}(p)\}$ (counted with multiplicity) is constant.*

Proof. Let γ be a smooth curve in D_2 connecting two points p and q and let U be a finite union of open discs compactly supported in D_1 which together cover the compact set $f^{-1}(\gamma)$. Since the image of any boundary component of U cannot cross γ the winding numbers of the image of any such boundary component with respect to p and q must be the same. Since that winding number counts the number of preimages inside that component we get by adding up the winding numbers for the different boundary components that $N_p = N_q$. □

Proof of Theorem 3.1. First assume that Φ is ω_{FS} -harmonic along a holomorphic disc $\{(g(\tau), \tau)\}$. We want to show that $g(\tau)$ must be of the form stated in the theorem.

Let $f(\tau) := \tau g(\tau)$. From Proposition 2.3 we have $\tilde{\Phi}$ is ω_{FS} -harmonic along the punctured disc $\{(f(\tau), \tau) : 0 < |\tau| < 1\}$ and since $\tilde{\Phi}$ is bounded away from $(0, 0)$ it will be ω_{FS} -harmonic along the unpunctured disc $\{(f(\tau), \tau) : |\tau| < 1\}$ unless that disc passes through $(0, 0)$. If it does pass through $(0, 0)$, the restriction of $\tilde{\Phi}$ to this disc has a logarithmic singularity of order at most one at $(0, 0)$, because $\tilde{\Phi}(z, \tau)$ is bounded from below by $\phi(z) + \ln |\tau|^2$.

Pick a τ_0 , $0 < |\tau_0| < 1$ and let $z_0 := f(\tau_0)$ and $t_0 := H(z_0, \tau_0)$. The restriction of

$$\psi_{t_0}(z) + (1 - t_0) \ln |\tau|^2 - \tilde{\Phi}(z, \tau)$$

to the disc is clearly subharmonic and less than or equal to zero and from Proposition 2.9 we see that it is equal to zero at (z_0, τ_0) . By the maximum principle we thus get that

$$\tilde{\Phi}(f(\tau), \tau) = \psi_{t_0}(f(\tau)) + (1 - t_0) \ln |\tau|^2 \quad (7)$$

for all τ .

We now consider the case when f is constant. Putting $f(\tau) = z_0$ in (7) we get that

$$\tilde{\Phi}(z_0, \tau) = \psi_{t_0}(z_0) + (1 - t_0) \ln |\tau|^2. \quad (8)$$

One possible case is that $z_0 = 0$, and in fact we will see later that $\tilde{\Phi}$ always is ω_{FS} -harmonic along this disc. If $z_0 \neq 0$, then we know that the LHS is ω_{FS} -harmonic along the whole unpunctured disc, which forces $t_0 = 1$ (otherwise the RHS would be singular at the center). Letting $\tau \rightarrow 1$ in (8) we get that

$$\psi_1(z_0) = \lim_{\tau \rightarrow 1} \tilde{\Phi}(z_0, \tau) = \phi(z_0)$$

by continuity of $\tilde{\Phi}$ which shows that $z_0 \in \Omega_1^c$.

We move on to the case when f is nonconstant. By (7) we then get that ψ_{t_0} is ω_{FS} -harmonic in a neighbourhood of any point $f(\tau)$, $|\tau| > 0$, which implies that $f(\tau) \in \Omega_{t_0}$. Note that this rules out the possibility of $t_0 = 0$, which in turn implies that $f(\tau) \neq 0$ whenever $\tau \neq 0$ as this would make the RHS in (7) be $-\infty$. If $f(0) \neq 0$ then $\tilde{\Phi}$ is ω_{FS} -harmonic along the entire disc and the same argument as above shows that $f(0) \in \Omega_{t_0}$ while $f(0) = 0$ of course also implies that $f(0) \in \Omega_{t_0}$. We thus have that f maps \mathbb{D} to Ω_t .

We now claim that f is proper. Because if we choose a sequence τ_i such that $|\tau_i| \rightarrow 1$ then by continuity of $\tilde{\Phi}$ and (7) we get that

$$\lim_{i \rightarrow \infty} (\psi_{t_0}(f(\tau_i)) - \phi(f(\tau_i))) = 0. \quad (9)$$

Since Ω_{t_0} is exhausted by the compact sets $\{z : \psi_{t_0}(z) \leq \phi(z) - 1/n\}$ by (9) $f(\tau_i)$ escapes to infinity in Ω_{t_0} . This shows that f is proper.

We now want to calculate $N_0 := \#\{f^{-1}(0)\}$. Because $f(\tau) \neq 0$ whenever $\tau \neq 0$, N_0 equals multiplicity of the zero at zero. If $f(0) \neq 0$ we have $N_0 = 0$, which by Lemma 3.2 is impossible, so we conclude that $f(0) = 0$. From (7) we get that

$$\psi_{t_0}(f(\tau)) + (1 - t_0) \ln |\tau|^2 = \tilde{\Phi}(f(\tau), \tau) \geq \phi(f(\tau)) + \ln |\tau|^2.$$

If m is the multiplicity of the zero of f at zero we get that the LHS has Lelong number $1 + (m - 1)t_0$ while the RHS has Lelong number one, which shows that $N_0 = m = 1$. This via Lemma 3.2 implies that f is a bijection between \mathbb{D} and Ω_{t_0} .

Since $\tau g(\tau) = f(\tau)$ this concludes the proof of the first direction of the Theorem.

We now prove that if $g(\tau)$ is of the form specified in the Theorem then Φ is ω_{FS} -harmonic along the graph of g . As before we let $f(\tau) := \tau g(\tau)$ and it is clearly enough to show that $\tilde{\Phi}$ is ω_{FS} -harmonic along the punctured disc $\{(f(\tau), \tau) : 0 < |\tau| < 1\}$.

The first case was $g \equiv 0$ and thus $f \equiv 0$. From (6) we see that $\tilde{\Phi}(0, \tau) = \phi(0) + \ln |\tau|^2$ and thus $\tilde{\Phi}$ is indeed ω_{FS} -harmonic along $\{(0, \tau) : 0 < |\tau| < 1\}$. The second case corresponded to $f(\tau) = z$ where $z \in \Omega_1^c$. From (6) we get that $\tilde{\Phi}(z, \tau) \geq \psi_1(z) = \phi(z)$. On the other hand the opposite inequality always holds and hence $\tilde{\Phi}$ is constant along the disc $\{(z, \tau) : |\tau| < 1\}$. The third case meant that f was a bijection between \mathbb{D} and some simply connected Hele-Shaw domain Ω_t , and $f(0) = 0$. From (6) we get that

$$\tilde{\Phi}(f(\tau), \tau) \geq \psi_t(f(\tau)) + (1 - t) \ln |\tau|^2.$$

The LHS is ω_{FS} -subharmonic with Lelong number one at $\tau = 0$ while the RHS is ω_{FS} -harmonic except at $\tau = 0$ where it also has Lelong number one. Since $f(\tau)$ escapes to

infinity in Ω_t as $|\tau| \rightarrow 1$ the RHS approaches the LHS as τ approaches the boundary. By the maximum principle we get the equality

$$\tilde{\Phi}(f(\tau), \tau) = \psi_t(f(\tau)) + (1 - t) \ln |\tau|^2,$$

which shows that $\tilde{\Phi}$ is ω_{FS} -harmonic along the punctured disc $\{(f(\tau), \tau) : 0 < |\tau| < 1\}$. This concludes the proof. \square

4. RESTATEMENT OF MAIN THEOREMS

We saw in Section 1.2 that for some ϕ the Hele-Shaw domains Ω_t are multiply connected for all t in some interval. Combining this with our main theorem yields the following.

Theorem 4.1. *Let ϕ be a Kähler potential on \mathbb{P}^1 whose Hele-Shaw flow satisfies the conclusion of Proposition 1.4, i.e. there are two times $0 < t_1 < t_2 < 1$ such that for any $t \in (t_1, t_2)$ Ω_t is multiply connected. Let Φ be the solution to the HMAE on $\mathbb{P}^1 \times \mathbb{D}$ with $\phi(z, \tau) := \phi(\rho(\tau)z)$ as boundary data. Then for such Φ there exists an open set U in $\mathbb{P}^1 \times \mathbb{D}$ with nonempty intersection with $\mathbb{P}^1 \times \partial\mathbb{D}$ which does not intersect any harmonic disc of Φ .*

Proof. By Theorem 3.1 no harmonic disc of Φ can intersect the open set

$$U := \{(z, \tau) : t_1 - 1 < H(\tau z, \tau) < t_2 - 1, |\tau| > 0\}.$$

Since $H(z, 1)$ clearly attains both values -1 and 0 it follows from continuity that $U \cap (\mathbb{P}^1 \times \partial\mathbb{D})$ is nonempty. \square

Theorem 4.2. *Let ϕ be as in Theorem 4.1. Then there exist a nonempty open set U' in $\mathbb{P}^1 \times \mathbb{D}$ and an $\epsilon > 0$ such that if $\phi'(z, \tau)$ is any smooth boundary data with*

$$\|\phi' - \phi\|_{C^2(\mathbb{P}^1 \times \partial\mathbb{D})} < \epsilon$$

and Φ' is the associated solution to the HMAE, then no harmonic disc of Φ' can pass through U' .

5. PROOF OF THEOREM 4.2

Recall from the Introduction that a solution Φ to the HMAE is called a *regular solution* if it is smooth and $\omega + dd^c\Phi(\cdot, \tau)$ is a Kähler form for all $\tau \in \mathbb{D}$. To prove Theorem 4.2 we will need Donaldson's Openness Theorem:

Theorem 5.1 (Donaldson [12, Theorem 1]). *The set of smooth boundary data $\phi : X \times \partial\mathbb{D} \rightarrow \mathbb{R}$ for which the HMAE has a regular solution is open in the C^2 topology.*

Proof of Theorem 4.2. Let U and Φ be as above, and U' be an nonempty open set which is relatively compact in $U \cap ((\mathbb{P}^1 \setminus \{0\}) \times \mathbb{D}^\times)$. We argue by contradiction, so assume that there is a sequence of solutions Φ_k with boundary values ϕ_k such that

$$\|\phi_k - \phi(z, \tau)\|_{C^2(\mathbb{P}^1 \times \partial\mathbb{D})} < 1/k$$

and such that for each k the graph of some holomorphic $g_k : \mathbb{D} \rightarrow \mathbb{P}^1$ is a harmonic disc of Φ_k which pass through U' .

Our first claim is that there exists an $r > 0$ such that for k large

$$\inf\{|g_k(\tau)| : \tau \in \mathbb{D}\} \geq r.$$

For simplicity we will assume that $\omega_\phi = \delta\omega_{FS}$ in $D_{r'} := r'\mathbb{D}$ for some $\delta, r' > 0$, as was the case for the ϕ used to prove Proposition 1.4. Then the function $h(z) := \phi(z) + (1 - \delta)\ln(1 + |z|^2)$ is harmonic in $D_{r'}$. By symmetry, it is not hard to see that the Hele-Shaw domains (defined by our potential ϕ) are initially just concentric discs centered at the origin, at least up to the point where $\Omega_t = D_{r'}$. From Theorem 3.1 we see that this implies that for any fixed $z \in D_{r'}$, the set $\{(z, \tau) : \tau \in \mathbb{D}\}$ is a harmonic disc associated to Φ . Thus $\Phi(z, \tau)$ is harmonic in τ for these fixed z .

This then implies that

$$\Phi(z, \tau) = -(1 - \delta)\ln(1 + |z|^2) + h(\tau z) \text{ on } D_{r'} \times \overline{\mathbb{D}}.$$

Pick a smooth Kähler potential u on \mathbb{P}^1 which is rotation invariant, and such that on $D_{r'}$ we have that

$$u \leq -(1 - \delta)\ln(1 + |z|^2)$$

with equality on $D_{r'/2}$ but strict inequality on $\partial D_{r'}$. Clearly $\Psi(z, \tau) := u(z)$ is a regular solution to the HMAE and we note that

$$\begin{aligned} \Psi(z, \tau) + h(\tau z) &= \Phi(z, \tau) \text{ on } D_{r'/2} \times \overline{\mathbb{D}}, \\ \Psi(z, \tau) + h(\tau z) &\leq \Phi(z, \tau) \text{ on } D_{r'} \times \overline{\mathbb{D}}, \\ \Psi(z, \tau) + h(\tau z) &< \Phi(z, \tau) \text{ on } \partial D_{r'} \times \overline{\mathbb{D}}. \end{aligned} \tag{10}$$

(Even if one loses the assumption that $\omega_\phi = \delta\omega_{FS}$ near 0 then arguing as in [20] one can still find an $r' > 0$ and harmonic h and regular solution Ψ such that the three statements of (10) hold).

Let $f_k(z, \tau) := \phi_k(z, \tau) - \phi(z, \tau)$ and let Ψ_k denote the solution to the HMAE with boundary data $\psi_k(z, \tau) := u(z) + f_k(z, \tau)$. Since by assumption

$$\|f_k\|_{C^2(\mathbb{P}^1 \times \partial\mathbb{D})} < 1/k$$

it follows from Donaldson's Openness Theorem that for large k the function Ψ_k is regular, and that the harmonic discs of Ψ_k converge to those of Ψ . In particular there exists an $r > 0$ such that for any large enough k there is a neighbourhood U_k of $D_r \times \mathbb{D}$ that is relatively compact in $D_{r'/2} \times \mathbb{D}$ which is foliated by harmonic discs of Ψ_k . We also have that for large k

$$\begin{aligned} \Psi_k(z, \tau) + h(\tau z) &= \Phi_k(z, \tau) \text{ on } D_{r'/2} \times \partial\mathbb{D}, \\ \Psi_k(z, \tau) + h(\tau z) &\leq \Phi_k(z, \tau) \text{ on } D_{r'} \times \partial\mathbb{D}, \\ \Psi_k(z, \tau) + h(\tau z) &< \Phi_k(z, \tau) \text{ on } \partial D_{r'} \times \overline{\mathbb{D}}, \end{aligned}$$

and therefore

$$\Psi_k(z, \tau) + h(\tau z) \leq \Phi_k(z, \tau) \text{ on } D_{r'} \times \overline{\mathbb{D}}.$$

On U_k though we must have

$$\Psi_k(z, \tau) + h(\tau z) \geq \Phi_k(z, \tau)$$

since along any harmonic disc of Ψ_k with boundary in $D_{r'/2} \times \partial\mathbb{D}$ the LHS is $\pi^*\omega_{FS}$ -harmonic, the RHS is $\pi^*\omega_{FS}$ -subharmonic, and they agree on the boundary. In conclusion

$$\Psi_k(z, \tau) + h(\tau z) = \Phi_k(z, \tau)$$

on U_k which shows Φ_k is regular on U_k and that U_k is foliated by harmonic discs of Φ_k .

By shrinking U' if necessary we may assume $U' \cap (D_{r'/2} \times \mathbb{D}) = \emptyset$, and so

$$U' \cap U_k = \emptyset \text{ for all } k.$$

Since the graph of g_k by assumption passes through U' it is not one of the harmonic discs that foliate U_k and since Φ_k is regular this means that the graph of g_k cannot intersect U_k . Finally this implies that $|g_k(\tau)| \geq r$ for all k as claimed.

Thus $1/g_k$ is a bounded family of holomorphic functions on \mathbb{D} so by Montel's theorem it is normal and thus replacing g_k by a subsequence we may assume g_k converges uniformly on compacts to a holomorphic function g . We claim that the graph of g then is a harmonic disc of Φ .

To see this let

$$h_k(\tau) := \Phi_k(g_k(\tau), \tau) + \ln(1 + |1/g_k(\tau)|^2)$$

and

$$h(\tau) := \Phi(g(\tau), \tau) + \ln(1 + |1/g(\tau)|^2).$$

By assumption h_k is harmonic and we want to show that h is harmonic (which says precisely that the graph of g is a harmonic disc of Φ). Now

$$|h_k(\tau) - h(\tau)| \leq |\Phi_k(g_k(\tau), \tau) - \Phi(g_k(\tau), \tau)| + |\Phi(g_k(\tau), \tau) - \Phi(g(\tau), \tau)|. \quad (11)$$

The first term of the RHS tends uniformly to zero since (as is easily seen)

$$\|\Phi_k - \Phi\|_{C^0(\mathbb{P}^1 \times \overline{\mathbb{D}})} \leq \|\phi_k - \phi\|_{C^0(\mathbb{P}^1 \times \partial\mathbb{D})} \leq \|\phi_k - \phi\|_{C^2(\mathbb{P}^1 \times \partial\mathbb{D})} \leq 1/k.$$

The second term of the RHS of (11) tends to zero uniformly on compacts since g_k tends to g uniformly on compacts and Φ is C^1 . Thus h_k tends to h uniformly on compacts which implies that h is harmonic, and so the graph of g is a harmonic disc of Φ . But since U' was compactly supported in $U \cap ((\mathbb{P}^1 \setminus \{0\}) \times \mathbb{D}^\times)$ the graph of g must intersect U which by Theorem 4.1 is a contradiction. This concludes the proof. \square

6. PARTIAL AND ALMOST SMOOTHNESS

Definition 6.1. Let Φ be upper semicontinuous on an open subset of $\mathbb{P}^1 \times \overline{\mathbb{D}}$ with $\pi_{\mathbb{P}^1}^* \omega_{FS} + dd^c \Phi \geq 0$ and $(\pi_{\mathbb{P}^1}^* \omega_{FS} + dd^c \Phi)^2 = 0$. The *regular locus* \mathcal{R}_Φ is the set of all $(z, \tau) \in \mathbb{P}^1 \times \overline{\mathbb{D}}$ near which Φ is smooth and $\pi_{\mathbb{P}^1}^* \omega_{FS} + dd^c \Phi(\cdot, \tau)$ is a Kähler form.

In particular the above definition applies to the weak solution of the HMAE. It is clear that \mathcal{R}_Φ is open, and inside it the kernel of $\pi_{\mathbb{P}^1}^* \omega_{FS} + dd^c \Phi$ defines a complex one-dimensional integrable distribution \mathcal{D}_Φ in \mathcal{R}_Φ . By restriction this gives a foliation on any open subset $V \subset \mathcal{R}_\Phi$ which, when necessary, we refer to as the *Monge-Ampère foliation* in V .

The following definition is a slight adaptation of those from [7, Sec 1.3].

Definition 6.2. We say the weak solution Φ to the HMAE is *partially smooth*¹ if it holds that (1) $\mathcal{R}_\Phi \cap (\mathbb{P}^1 \times \mathbb{D})$ is foliated by harmonic discs and (2) \mathcal{R}_Φ is dense in $\mathbb{P}^1 \times \partial\mathbb{D}$ and (3) the fibrewise volume form $\omega_{\Phi(\cdot, \tau)}$ which is defined on R_ϕ extends to a continuous $(1, 1)$ -form on $\mathbb{P}^1 \times \mathbb{D}$.

The following should be compared with [7, Thm. 1.3.2] which says weak solutions are always partially smooth.

Corollary 6.3. *There exist ϕ such that the weak solution Φ to the HMAE with boundary data $\phi(z, \tau) := \phi(\rho(\tau)z)$ are not partially smooth.*

¹Prof Tian has informed us that the definition of partially smooth in [7, Def 1.3.1] contains a typo, and should read that “ R_ϕ is saturated in $X \times (\Sigma \setminus \partial\Sigma)$ ”. This is potentially weaker than what is written in [7, Def 1.3.1], and is what we use here.

Proof. This is immediate, for our example from Theorem 4.1 gives boundary data for which the solution to the HMAE is not partially smooth, since no harmonic disc intersects the open set U which has non-empty intersection with the boundary. \square

Remark 6.4. The example in Theorem 4.2 is in apparent contradiction with [7, Thm 1.3.4], since the solutions Φ' associated to the perturbed boundary data ϕ' cannot be “almost smooth” (for by [7, Prop 2.3.1] for any almost smooth solution, the set of harmonic discs is dense in $\mathbb{P}^1 \times \mathbb{D}$).

REFERENCES

- [1] E Bedford and D Burns *Holomorphic mapping of annuli in \mathbb{C}^n and the associated extremal function*. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 6 (1979), no. 3, 381–414.
- [2] R Berman *Bergman kernels and equilibrium measures for ample line bundles* (2007) Preprint arXiv:0704.1640.
- [3] R Berman *On the optimal regularity of weak geodesics in the space of metrics on a polarized manifold* (2014) Preprint arXiv:1405.6482.
- [4] R Berman and B Berndtsson *Convexity of the K-energy on the space of Kähler metrics* (2014) Preprint arXiv:1405.0401
- [5] Z Błocki *On geodesics in the space of Kähler metrics* Advances in Geometric Analysis, ed. S. Janeczko et al., Advanced Lectures in Mathematics 21, pp. 3–20, International Press, 2012.
- [6] X X Chen *The Space of Kähler Metrics* J. Differential Geom. Volume 56, Number 2 (2000), 189–234.
- [7] X X Chen and G Tian *Geometry of Kähler metrics and foliations by holomorphic discs*. Publ. Math. Inst. Hautes Études Sci. No. 107 (2008), 1–107.
- [8] L.A. Caffarelli and N. M. Rivière *Smoothness and analyticity of free boundaries in variational inequalities* Annali della Scuola Normale Superiore di Pisa, Classe di Scienze 4em série, tome 3, no 2 (1976), p. 289–310.
- [9] L.A. Caffarelli and D. Kinderlehrer *Potential methods in variational inequalities* J. Analyse Math. 37 (1980), 285–295.
- [10] T Darvas and L Lempert *Weak geodesics in the space of Kähler metrics* Math. Res. Lett. 19 (2012), no. 5, 1127–1135.
- [11] T Darvas *Morse theory and geodesics in the space of Kähler metrics* Proc. Amer. Math. Soc. 142 (2014), no. 8, 2775–2782.
- [12] S K Donaldson *Holomorphic discs and the complex Monge-Ampère equation*. J. Symplectic Geom. 1 (2002), no. 2, 171–196.
- [13] S K Donaldson *Nahm’s equations and free-boundary problems*. The many facets of geometry, 71–91, Oxford Univ. Press, Oxford, 2010.
- [14] Vincent Guedj (Editor) *Complex Monge-Ampère Equations and Geodesics in the Space of Kähler Metrics* Lecture Notes in Mathematics 2038, Springer 2012
- [15] B. Gustafsson and A. Vasil’ev *Conformal and potential analysis in Hele-Shaw cells* Advances in Mathematical Fluid Fluid Mechanics. Birkhäuser Verlag, Basel, 2006.
- [16] H. Hedenmalm and S. Shimorin *Hele-Shaw flow on hyperbolic surfaces* J. Math. Pures Appl. (9) 81 (2002), no. 3, 187–222.
- [17] C Kiselman *The partial Legendre transformation for plurisubharmonic functions* Invent. Math. 49 (1978), 137–148.
- [18] L Lempert and L Vivas *Geodesics in the space of Kähler metrics* Duke Math. J. 162 (2013), no. 7, 1369–1381.
- [19] R. T. Rockafellar *Convex analysis* Princeton Mathematical Series, No. 28, Princeton University Press, Princeton, N.J. 1970.
- [20] J Ross and D Witt Nyström *The Hele-Shaw flow and moduli of holomorphic discs* (2012) Preprint arXiv:1212.2337.
- [21] S Semmes *Complex Monge-Ampère and symplectic manifolds* Amer. J. Math. 114 (1992), no. 3, 495–550.

JULIUS ROSS, DPMMS, UNIVERSITY OF CAMBRIDGE, UK. J.ROSS@DPMMS.CAM.AC.UK

DAVID WITT NYSTRÖM, DPMMS, UNIVERSITY OF CAMBRIDGE, UK.
D.WITTYNYSTROM@DPMMS.CAM.AC.UK, DANSPOLITIK@GMAIL.COM